

The Signature of the Symmetric Union of Links

FURUTA Yuuichi and SHINOHARA Yaichi

§ 1. Introduction

Let L be an unoriented link in R^3 and $D(L)$ a link diagram of L on R^2 . First we assume that $D(L)$ is connected. R^2 is divided into connected regions by $D(L)$ which can be alternately colored white and black in such a way that adjacent regions are of distinct colors and the unbounded region is a white region. For each crossing P of $D(L)$, if the white regions at P are distinct, we assign the incident number $\theta(P)$ with 1 or -1 depending on whether the overcrossing line coincides with the undercrossing line by turning over the black regions at P counterclockwise or clockwise. If the white regions at P are same, then we define $\theta(P)$ to be 0. Let $W_0, W_1, W_2, \dots, W_m$ be the white regions. For $0 \leq i \leq m$ and $0 \leq j \leq m$, we define g_{ij} as follows:

$$g_{ij} = \begin{cases} \sum_{P \in \partial W_i} \theta(P) & \text{if } i = j, \\ - \sum_{P \in W_i \cap W_j} \theta(P) & \text{if } i \neq j. \end{cases}$$

Define the $(m+1) \times (m+1)$ matrix $\tilde{G}(L)$ by $\tilde{G}(L) = (g_{ij})$. The $m \times m$ matrix $G(L)$ obtained from $\tilde{G}(L)$ by deleting the row and the column corresponding to any white region W_{i_0} is called *the Goeritz matrix associated to $D(L)$* (see [2], [4], [5] and [11]).

If $D(L)$ is disconnected, then $D(L)$ can be expressed as the union of a finite number of link diagrams $D(L_1), D(L_2), \dots, D(L_n)$ such that $D(L_1), D(L_2), \dots, D(L_n)$ are connected and are disjoint of each other. We construct $\tilde{G}(L_1), \tilde{G}(L_2), \dots, \tilde{G}(L_n)$ for link diagrams $D(L_1), D(L_2), \dots, D(L_n)$, respectively. Then we define the matrix $\tilde{G}(L)$ by

$$\tilde{G}(L) = \begin{pmatrix} \tilde{G}(L_1) & 0 & \cdots & 0 \\ 0 & \tilde{G}(L_2) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \tilde{G}(L_n) \end{pmatrix}.$$

The matrix $G(L)$ obtained from $\tilde{G}(L)$ by deleting the i_0 -th row and the i_0 -th column for any i_0 is called *the Goeritz matrix associated to $D(L)$* for this case.

Now let L be an oriented link and $D(L)$ its diagram. A crossing P of $D(L)$ is said to be *of type II* if the oriented overcrossing line coincides with the oriented undercrossing line by turning over the black regions at P . Otherwise, it is said to be *of type I*. We define $\mu(D(L)) = \sum \theta(P)$ (summed over all crossings P of type II).

For a symmetric matrix A , $\sigma(A)$ will denote the signature of A , i.e., when A is diagonalized, the signature of A is defined to be the number of positive diagonal elements minus the number of negative diagonal elements (see [8], [10]). It is well-known that for an oriented link L , $\sigma(G(L)) - \mu(D(L))$ is an invariant of the oriented link type of L , which is denoted by $\sigma(L)$ and is called *the signature of an oriented link L* (see[4], [6]). In [4], Y. Furuta and Y. Shinohara showed that for an oriented link L and its diagram $D(L)$, $\chi(L) = \text{rank}G(L) - \mu(D(L)) \pmod{2}$ is an invariant of the oriented link type of L .

Let L be a link in R^3 . As shown in Fig. 1.1, first we take a connected sum of L and its mirror image L^* , and secondly we twist two corresponding parallel arcs, one from L and the other from L^* , n times. Here, the twisting number n is defined to be positive or negative according as shown in Fig. 1.2. The resulting link is denoted by $S(L, n)$ and is called *a symmetric union* if n is even and *a skew-symmetric union* if n is odd.




Fig. 1.1




Fig. 1.2

The symmetric union and the skew-symmetric union were introduced by S. Kinoshita and H. Terasaka in [9]. For the case that L is a trivial knot, R. Fukumoto and Y. Shinohara computed the Lickorish-Millett polynomial for $S(L, 2n)$, by using which they classified $S(L, 2n)$ in [3]. In this paper, we will study the signature of $S(L, n)$ and obtain the following theorem:

THEOREM 1.1 *If $S(L, n)$ is a symmetric union of an oriented link L , i.e., n is even, then we have*

$$\sigma(S(L, n)) = \begin{cases} 0 & \text{if } \chi(S(L, n)) \equiv 0 \pmod{2}, \\ -\epsilon & \text{if } \chi(S(L, n)) \equiv 1 \pmod{2}, \end{cases}$$


where ϵ denotes the sign of n .

We also give an alternate proof for the following theorem which was proved by S. Asakura and Y. Shinohara [1]:

THEOREM 1.2 *If n is even, then for any oriented link L with $\det L \neq 0$ we have $\sigma(S(L, n)) = 0$.*

§ 2. The Signature of the Goeritz Matrix of $S(L, n)$

The link diagram $D(S(L, n))$ of $S(L, n)$ is given in Fig. 2.1.

**Fig. 2.1**

The Goeritz matrix $G(S(L, n))$ associated to $D(S(L, n))$ has the following form:

$$G(S(L, n)) = \begin{pmatrix} G(L) & 0 & 0 & {}^tX \\ 0 & -G(L) & 0 & -{}^tX \\ 0 & 0 & T_{|n|-1} & {}^tY_{|n|-1} \\ X & -X & Y_{|n|-1} & \epsilon \end{pmatrix}.$$

Here, $G(S(L, n))$ is obtained from $\tilde{G}(S(L, n))$ by deleting the row and the column corresponding to the white region W_0 , and the last row and the last column of $G(S(L, n))$ correspond to the white region $W_{|n|''}$. Also, $G(L)$ is the Goeritz matrix for L which is an $m \times m$ matrix, ϵ denotes the sign of n and X is a $1 \times m$ matrix. For a positive integer k , Y_k is the $1 \times k$ matrix whose $(1, j)$ -th entry is $-\epsilon$ if $j = k$ and is 0 otherwise, and T_k is the $k \times k$ matrix given by

$$T_k = \begin{pmatrix} 2\epsilon & -\epsilon & 0 & 0 & \cdots & 0 \\ -\epsilon & 2\epsilon & -\epsilon & 0 & \cdots & 0 \\ 0 & -\epsilon & 2\epsilon & -\epsilon & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & -\epsilon & 2\epsilon & -\epsilon \\ 0 & 0 & \cdots & 0 & -\epsilon & 2\epsilon \end{pmatrix}.$$

First we show the following lemma:

LEMMA 2.1 For a positive integer k , we have $\det T_k = (k+1) \epsilon^k$ and $\sigma(T_k) = k\epsilon$.

Proof. By expanding the first row of T_k , we obtain the following recursion formula:

$$\det T_k = 2\epsilon \det T_{k-1} - \det T_{k-2} \quad (k \geq 3),$$

which implies that $\det T_k = (k+1) \epsilon^k$. Since

$$\Delta_0 = 1, \Delta_1 = \det T_1, \Delta_2 = \det T_2, \dots, \Delta_k = \det T_k$$

is a σ -series for T_k , we have

$$\sigma(T_k) = \sum_{i=0}^{k-1} \text{sign}(\Delta_i \cdot \Delta_{i+1}) = \sum_{i=0}^{k-1} \epsilon = k\epsilon.$$

(For a σ -series, see [8], [10].)

Q.E.D

Now we prove the following lemma which plays a crucial role in the proof of our theorems:

LEMMA 2.2

- (1) If $\text{rank}(G(L) \setminus X) = \text{rank} G(L) + 1$, then
 $\text{rank} G(S(L, n)) = 2\text{rank} G(L) + |n| + 1$ and $\sigma(G(S(L, n))) = (|n| - 1) \epsilon$.
- (2) If $\text{rank}(G(L) \setminus X) = \text{rank} G(L)$, then
 $\text{rank} G(S(L, n)) = 2\text{rank} G(L) + |n|$ and $\sigma(G(S(L, n))) = |n| \epsilon = n$.

Proof. First we consider the case that $\text{rank}(G(L) \setminus X) = \text{rank} G(L) + 1$. Since $G(L)$ is a symmetric matrix, there exists an orthogonal matrix P such that

$${}^t P G(L) P = \begin{pmatrix} \lambda_1 & & & & & \\ & \lambda_2 & & & & \\ & & \ddots & & & \\ & & & \lambda_r & & \\ & & & & \lambda_{r+1} & \\ & 0 & & & & \ddots \\ & & & & & & \lambda_m \end{pmatrix},$$

where $\lambda_1, \lambda_2, \dots, \lambda_r$ are non-zero real numbers and $\lambda_{r+1}, \dots, \lambda_m$ are 0. Note that $r = \text{rank} G(L)$.

Let

$$Q = \begin{pmatrix} P & 0 & 0 & 0 \\ 0 & P & 0 & 0 \\ 0 & 0 & E_{|n|-1} & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Then

$${}^tQA(S(L, n))Q = \begin{pmatrix} {}^tPG(L)P & 0 & 0 & {}^t(XP) \\ 0 & -{}^tPG(L)P & 0 & {}^t(-XP) \\ 0 & 0 & T_{|n|-1} & {}^tY_{|n|-1} \\ XP & -XP & Y_{|n|-1} & \epsilon \end{pmatrix}.$$

Since

$${}^tP(G(L) \ {}^tX) \begin{pmatrix} P & 0 \\ 0 & 1 \end{pmatrix} = ({}^tPG(L)P \ {}^t(XP)),$$

it follows that

$$\text{rank}({}^tPG(L)P \ {}^t(XP)) = \text{rank}(G(L) \ {}^tX) = \text{rank} G(L) + 1 = \text{rank}({}^tPG(L)P) + 1.$$

If we put $XP = (x_1, x_2, \dots, x_r, x_{r+1}, \dots, x_m)$, then there exists an integer r_0 ($r+1 \leq r_0 \leq m$) such that x_{r_0} is not zero. Hence, there exists a non-singular matrix R such that

$${}^tRG(S(L, n))R = \begin{pmatrix} A_r & 0 & 0 & 0 & 0 \\ 0 & -A_r & 0 & 0 & 0 \\ 0 & 0 & T_{|n|-1} & 0 & 0 \\ 0 & 0 & 0 & B & 0 \\ 0 & 0 & 0 & 0 & C \end{pmatrix},$$

where

$$A_r = \begin{pmatrix} \lambda_1 & & & 0 \\ & \lambda_2 & & \\ & & \ddots & \\ 0 & & & \lambda_r \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

and C is the $(2m - 2r - 1) \times (2m - 2r - 1)$ zero matrix. Therefore we have

$$\text{rank}G(S(L, n)) = \text{rank}({}^tRG(S(L, n))R) = 2\text{rank}G(L) + |n| + 1.$$

Moreover, by using Lemma 2.1, we obtain

$$\sigma(G(S(L, n))) = \sigma({}^tRG(S(L, n))R) = r - r + \sigma(T_{|n|-1}) + \sigma(B) = (|n| - 1)\epsilon.$$

Next we consider the second case. Since $\text{rank}(G(L) {}^tX) = \text{rank}G(L)$, it follows from Theorem 7 in Chapter II of [7] that there exists a $1 \times m$ matrix Z such that $G(L) {}^tZ = {}^tX$. If we put

$$U = \begin{pmatrix} E_m & 0 & 0 & -{}^tZ \\ 0 & E_m & 0 & -{}^tZ \\ 0 & 0 & E_{|n|-1} & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

then

$${}^tUG(S(L, n))U = \begin{pmatrix} G(L) & 0 & 0 & 0 \\ 0 & -G(L) & 0 & 0 \\ 0 & 0 & T_{|n|-1} & {}^tY_{|n|-1} \\ 0 & 0 & Y_{|n|-1} & \epsilon \end{pmatrix}.$$

Let $V = \begin{pmatrix} T_{|n|-1} & {}^tY_{|n|-1} \\ Y_{|n|-1} & \epsilon \end{pmatrix}$. Then by using Lemma 2.1 we obtain

$$\det V = \epsilon \det T_{|n|-1} + \epsilon(-\epsilon) \det T_{|n|-2} = \epsilon^{|n|} = \epsilon^n.$$

This implies that

$$\text{rank}G(S(L, n)) = 2\text{rank}G(L) + |n|.$$

Since

$$\Delta_0 = 1, \Delta_1 = \det T_1, \Delta_2 = \det T_2, \dots, \Delta_{|n|-1} = \det T_{|n|-1}, \Delta_{|n|} = \det V$$

is a σ -series for V , we have

$$\sigma(V) = \sum_{i=0}^{|n|} \text{sign}(\Delta_i \cdot \Delta_{i+1}) = \sum_{i=0}^{|n|} \epsilon = |n| \epsilon = n.$$

Hence we obtain

$$\sigma(G(S(L, n))) = \sigma(G(L)) - \sigma(G(L)) + \sigma(V) = n.$$

Q.E.D

§ 3. The Proof for Theorem 1.1 and Theorem 1.2.

In this section, we will prove Theorem 1.1 and Theorem 1.2. Since n is even, for an oriented link L , $S(L, n)$ will be shown in Fig. 3.1.




Fig. 3.1

It is easy to show that $\mu(D(S(L, n))) = n$.

First of all, we will prove Theorem 1.1. If $\chi(S(L, n)) \equiv 0 \pmod{2}$, then

$$\text{rank}G(S(L, n)) \equiv \mu(D(S(L, n))) = n \equiv 0 \pmod{2}.$$

Lemma 2.2 implies that $\text{rank}(G(L) \cdot X) = \text{rank}G(L)$. Thus we have

$$\sigma(G(S(L, n))) = |n| \epsilon = n \quad \text{and} \quad \sigma(S(L, n)) = n - n = 0.$$

If $\chi(S(L, n)) \equiv 1 \pmod{2}$, then

$$\text{rank}G(S(L, n)) \equiv \mu(D(S(L, n))) + 1 = n + 1 \equiv 1 \pmod{2}.$$

Again, by Lemma 2.2, we have $\text{rank}(G(L) \cup X) = \text{rank} G(L) + 1$, which yields

$$\sigma(G(S(L, n))) = |n| \epsilon - \epsilon \quad \text{and} \quad \sigma(S(L, n)) = n - \epsilon - n = -\epsilon.$$

This completes the proof for Theorem 1.1.

Finally, we will prove Theorem 1.2. Since $\det L = |\det G(L)| \neq 0$, $\text{rank}(G(L) \cup X) = \text{rank}(G(L))$. Hence, Lemma 2.2 implies that

$$\sigma(S(L, n)) = \sigma(S(L, n)) - \mu(D(S(L, n))) = n - n = 0,$$

which completes the proof for Theorem 1.2.

Reference

1. S. Asakura and Y. Shinohara, On the signature of the symmetric union and the skewsymmetric union of links, The Kwansei Gakuin University Natural Sciences Reviews, 10 (2006), 1-9.
2. G. Burde and H. Zieschang, *Knots*, Walter de Gruyter, 1985.
3. L. Fukumoto and Y. Shinohara, On the symmetric union of a trivial knot, Kobe J. Math., 14(1997), 123-132.
4. Y. Furuta and Y. Shinohara, On the Goeritz matrix of links, The Kwansei Gakuin University Natural Sciences Reviews, 12 (2008), 1-18.
5. L. Goeritz, Knoten und quadratische Formen, Math. Z. 36(1933), 647-654.
6. C.McA. Gordon and R.A. Litherland, On the signature of a link, *Inventiones mathematicae* 47(1978), 53-69.
7. N. Jacobson, *Lectures in Abstract Algebra*, Volume II, Van Nostrand Reinhold Company, 1953.
8. B.W.Jones, *The arithmetic theory of quadratic forms*, The Carus Mathematical Monographs, Buffalo, 1950.
9. S. Kinoshita and H. Terasaka, On unions of knots, Osaka Math. J. 9(1957), 131-153.
10. K. Murasugi, On a certain numerical invariant of link types, Trans. Amer. Math. Soc. 117(1965), 387-422.
11. K. Reidemeister, *Knotentheorie*, Berlin, Springer, 1932.